



Technical Data

10.99

Weight/Gallon
@ 68°F

1.318

Specific Gravity

7.7-8.5

pH @ 68°F

7° F

Freezing Point

DIRECTIONS FOR USE

When applied according to specified guidelines, accessS® effectively provides the recommended sulfur needs of most crops..

accessS can be applied:

- with with planter placement in a band away from the seed (ex: 2"x 2")
- broadcast or banded over the top of the seed zone
- as a topdress
- as a sidedress
- through fertigation

USE RATE SUMMARY TABLE

At Planting Application Rates

Gallons Per Acre

Field and Row Crops	0 - 20
Vegetables and Fruit Crops	0 - 10*
Orchards and Vineyards	0 - 10*

In-Season Application Rates - Per Application

Field and Row Crops	0 - 20	Sidedress or Fertigation
Vegetables and Fruit Crops	0 - 10	Sidedress or Fertigation
Orchards and Vineyards	0 - 10	Soil Application or Fertigation

Foliar Application Rates - Per Application

Field and Row Crops	Not Recommended
Vegetables and Fruit Crops	Not Recommended
Orchards and Vineyards	Not Recommended

* = Not recommended in transplant solution.

Composition Guaranteed Analysis

Total Nitrogen (N)

7.00%

7.00% Ammoniacal Nitrogen

Sulfur (S)

17.00%

17.00% Combined Sulfur

Iron (Fe)

0.25%

0.25% Water Soluble Iron (Fe)

Manganese (Mn)

0.05%

0.05% Water Soluble Manganese (Mn)

Zinc (Zn)

0.05%

0.05% Water Soluble Zinc (Zn)

Derived from: Ammonium Sulfate, Ferrous Sulfate,
Manganese Sulfate, Sulfuric Acid, Zinc Sulfate



Guaranteed by: AgroLiquid
Division of COG Marketers, Ltd.
3055 W M-21
St. Johns, MI 48879
agroliquid.com



Directions For Use General Guideline:

For proper agronomic application rates suitable for your geographical area or the maximum allowable non-nutrient application rate per acre, consult a trained soil specialist at AgroLiquid or call or write to AgroLiquid with the address provided.

accessS is a proprietary high-sulfur formulation intended for application anywhere five or more pounds of elemental sulfur is recommended. These enhanced efficiencies make accessS both environmentally and economically responsible.

Crop	In-Furrow
Wheat	1-3 gal/A

Broadcast, or banded not less than 2" from the seed furrow, surface banded, or applied through drip irrigation at the base of the plant		
RATE: 1-100 gallon/acre		
Corn	Cotton	Potato
Soybean	Sugarbeet	Alfalfa
Sorghum	Canola	
Dry Beans	Wheat	
RATE: 1-30 gallon/acre		
Grapes	Apples	Tree Fruit
Tomato	Tree Nuts	Vegetables
Tobacco		

In-Season Soil Application Recommendations RATE: 1-100 gallon/acre

Corn Sidedress or fertigation	Grapes Banded or through drip irrigation
Soybean Sidedress or fertigation	Tomato Banded or through drip irrigation
Sorghum Sidedress or fertigation	Tobacco Banded or through drip irrigation
Dry Beans Sidedress or fertigation	Tree Fruit Banded or through drip irrigation
Cotton Sidedress or fertigation	Tree Nuts Banded or through drip irrigation
Sugarbeet Sidedress or fertigation	Vegetables Banded or through drip irrigation
Canola Topdress or fertigation	
Wheat Topdress or fertigation	
Potato Sidedress or fertigation	
Alfalfa Sidedress or fertigation	

Please consult with an AgroLiquid Sales Account Manager or Agronomist for further direction when utilizing rates higher than the lower limit of the given range.

 **Guaranteed by: AgroLiquid**
 Division of COG Marketers, Ltd.
 3055 W M-21
 St. Johns, MI 48879
 AGROLIQUID. agroliquid.com

NOTE: Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>



GS07
WARNING - Irritant: Eyes, Skin