

Technical Data



DIRECTIONS FOR USE

The quality and composition of springuP[™] allows placement directly into the root zone with little risk of tissue damage, giving springuP greater efficacy and more application flexibility.

SpringuP can be applied:

- with versatile planter placement options
- banded over the top of the seed zone
- through fertigation
- with crop protection products

USE RATE SUMMARY TABLE

At Planting Application Rates	Gallons Per Acre
Field and Row Crops	0 - 20
Vegetables and Fruit Crops	0 - 20 or 2% in Transplant Solution
Orchards and Vineyards	0 - 20 or 2% in Transplant Solution

In-Season Application Rates - Per Application

Field and Row Crops	1 - 20	Sidedress or Fertigation
Vegetables and Fruit Crops	1 - 20	Sidedress or Fertigation
Orchards and Vineyards	1 - 20	Soil Application or Fertigation

Composition Guaranteed Analysis

Total Nitrogen (N) 4.00%

2.50% Ammoniacal Nitrogen 1.50% Urea Nitrogen

Available Phosphate (P₂O₅) 15.00%

> Soluble Potash (K_2O) 3.00%

Derived from: Ammonium Phosphate, Superphosphoric Acid, Potassium Carbonate, Urea



springduP

Directions For Use General Guideline:

For proper agronomic application rates suitable for your geographical area or the maximum allowable non-nutrient application rate per acre, consult a trained soil specialist at AgroLiquid or call or write to AgroLiquid with the address provided.

SpringuP[™] is an ideal starter liquid phosphorus solution for strong early-season performance because of its fast-release P and easy application in-furrow.

Сгор	In-Furrow
Corn (Grain) 30" Row Spacing	2-10 gal/A
Corn (Silage) 30" Row Spacing	2-10 gal/A
Soybeans 30" Row Spacing	1-3 gal/A
Soybeans 15" Row Spacing	2-7 gal/A
Sorghum	1-3 gal/A
Dry Beans	0-3* gal/A
Cotton	0-5* gal/A
Sugarbeet	0-3* gal/A
Canola Drilled	2-5 gal/A
Wheat (Spring or Winter)	2-10 gal/A
Potato	2-10 gal/A Direct contact with the seed piece
Alfalfa	1-3 gal/A

In-Season Soil Application Recommendations RATE: 1-20 gallon/acre

Corn Sidedress

Soybean 30" Rows Sidedress

Cotton Sidedress

Sugarbeet Sidedress

Wheat Topdress application up to growth stage Feekes 4

Potato Sidedress

Alflafa Topdress Broadcast, or banded not less than 2" from the seed furrow, surface banded, or applied through drip irrigation at the base of the plant RATE: 1-20 gallon/acre

Corn Soybean Sorghum	Canola Wheat Potato	Tobacco Apples Tree Nuts
Dry Beans	Alfalfa	Tree Fruit
Cotton Sugarbeet	Grapes Tomato	Vegetables

2% in transplant solution during transplanting.

Please consult with an AgroLiquid Sales Account Manager or Agronomist for further direction when utilizing rates higher than the lower limit of the given range.



NOTE: Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

