






Micro 600™

Technical Data



DIRECTIONS FOR USE

Micro 600™ may be used as a micronutrient component in any fertilizer application method. Each crop has minimum requirements for micronutrients in specific proportions to each other. The synergy of applying the combination of the nutrients found in Micro600™ benefits most cropping programs and soil types - especially in high pH soil conditions.

Micro 600™ can be applied:

- with versatile planter placement options
- as a sidedress
- as a foliar application
- through fertigation

Use Rate Summary Table - Micro 600

At Planting Application Rates	Gallons Per Acre
Field and Row Crops	0 - 2
Vegetables and Fruit Crops	0 - 2 or 0.25% in Transplant Solution
Orchards and Vineyards	0 - 2 or 0.25% in Transplant Solution
In-Season Application Rates - Per Application	
Field and Row Crops	0.125 - 2 Sidedress or Fertigation
Vegetables and Fruit Crops	0.125 - 2 Sidedress or Fertigation
Orchards and Vineyards	0.125 - 2 Soil Applied or Fertigation
Foliar Application Rates - Per Application	
Field and Row Crops	0.125 - 0.5
Vegetables and Fruit Crops	0.125 - 0.5
Orchards and Vineyards	0.125 - 0.5

Composition Guaranteed Analysis

Sulfur (S)
2.00%
2.00% Combined Sulfur

Boron (B)
0.02%

Copper (Cu)
0.05%
0.05% Water Soluble Copper (Cu)

Iron (Fe)
2.00%
2.00% Water Soluble Iron (Fe)

Manganese (Mn)
1.00%
1.00% Water Soluble Manganese (Mn)

Zinc (Zn)
1.00%
1.00% Water Soluble Zinc (Zn)

Derived from: Boric Acid, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, Zinc Sulfate

Warning: Contains boron. Do not use on boron-sensitive crops. Use only according to the directions given by a trained AgroLiquid soil specialist.



Guaranteed by: AgroLiquid
Division of COG Marketers, Ltd.
3055 W M-21
St. Johns, MI 48879
agroliquid.com



Directions For Use General Guideline:

For proper agronomic application rates suitable for your geographical area or the maximum allowable non-nutrient application rate per acre, consult a trained soil specialist at AgroLiquid or call or write to AgroLiquid with the address provided.

The Flavonol Polymer Technology in Micro 600™ allows for effective uptake and utilization of the micronutrients by the plant. Applications at planting, as foliar, or through irrigation all produce excellent results. This product can be used as a recovery aid or as a preventative measure to overcome the symptoms of iron chlorosis – even in soils where soils tests indicate sufficient levels of iron.

Crop	In-Furrow
Corn (Grain)	0-1 gal/A
30" Row Spacing	
Corn (Silage)	0-1 gal/A
30" Row Spacing	
Soybeans	0-0.5 gal/A
30" Row Spacing	
Soybeans	0-0.5 gal/A
15" Row Spacing	
Sorghum	0-0.5 gal/A
Dry Beans	0-0.25* gal/A
Cotton	0-0.25* gal/A
Sugarbeet	0-0.25* gal/A
Canola	0-0.25* gal/A
Wheat	0-0.25* gal/A
(Spring or Winter)	
Potato	0-1 gal/A
	Direct contact with the seed piece
Alfalfa	0-0.25* gal/A

In-Season Soil Application

RATE: 0.125 - 2 gal/A unless otherwise noted.

Corn

Sidedress

Sorghum

Sidedress

Cotton

Sidedress

Sugarbeet

Sidedress

Wheat

Topdress up to Feekes Stage 4

Potato

Sidedress or fertigation

Alfalfa

Prior to, or within 14 days of spring green-up, and/or 0-7 days after cutting, broadcast

Grapes

Broadcast, surface banded or through drip irrigation at bud break or during the growing season

Tomato

Banded or through drip irrigation during the growing season

Tobacco

Banded or through drip irrigation during the growing season

Apples

Banded or through drip irrigation during the growing season

Tree Nuts

Banded or through drip irrigation during the growing season

Other Tree Fruits

Banded or through drip irrigation during the growing season

Vegetables

Broadcast, surface banded or through drip irrigation during the growing season

Foliar Application Recommendations

RATE: 0.125 -0.5 gal/A unless otherwise noted

Corn

Soybean 30" and 15" Rows

Sorghum

Dry Beans

Cotton

Sugarbeet

Canola

Wheat

Potato

Alfalfa

Grapes

Tomato

Tobacco

Apples

Tree Nuts

Other Tree Fruits

Vegetables

Broadcast, or banded not less than 2" from the seed furrow, surface banded, or applied through drip irrigation at the base of the plant

RATE: 0.125 -2 gal/A

Corn	Canola	Tobacco
Soybean	Wheat	Apples
Sorghum	Potato	Tree Nuts
Dry Beans	Alfalfa	Tree Fruit
Cotton	Grapes	Vegetables
Sugarbeet	Tomato	

0.25% in Transplant Solution

Grapes	Apples	Vegetables
Tomato	Tree Nuts	
Tobacco	Tree Fruit	

