




Technical Data



DIRECTIONS FOR USE

PrimAgro K can be used in virtually any cropping situation where potassium is recommended. PrimAgro K should be used under normal planting conditions, where soil temperatures permit proper germination.

PrimAgro K can be applied:

- with versatile planter placement options
- broadcast or banded over the top of the seed zone
- as a topdress
- as a sidedress
- through fertigation

In general, where liquid fertilizers such as PrimAgro K are applied in horticultural crop production, it is customary to split the seasonal fertilizer needs into a number of smaller applications. In this case PrimAgro K may be used for early-season potassium, while AgroLiquid Sure-K® may be more appropriate for mid- and late-season application. PrimAgro K is not recommended as a foliar.

Use Rate Summary Table

At Planting Application Rates		Gallons Per Acre
Field and Row Crops		0 - 20
Vegetables and Fruit Crops		0 - 10
Orchards and Vineyards		Not Recommended
In-Season Application Rates - Per Application		
Field and Row Crops	1 - 20	Sidedress or Fertigation
Vegetables and Fruit Crops	1 - 10	Sidedress or Fertigation
Orchards and Vineyards	1 - 8	Soil Application or Fertigation
Foliar Application Rates - Per Application		
Field and Row Crops	0 - 3*	
Vegetables and Fruit Crops	Not Recommended	
Orchards and Vineyards	Not Recommended	

* = Please consult with an AgroLiquid Sales Account Manager or Agronomist for further direction when utilizing rates higher than the lower limit of the given range.

Composition Guaranteed Analysis

Total Nitrogen (N)

1.00%

0.50% Nitrate Nitrogen
0.50% Ammoniacal Nitrogen

Soluble Potash (K₂O)

8.00%

Sulfur (S)

6.00%

6.00% Combined Sulfur

Derived from: Ammonium Nitrate,
Potassium Carbonate, Potassium Sulfate

ALSO CONTAINS NON PLANT FOOD INGREDIENTS

Contains 401 million total
colony forming units per gallon
(4.01x10⁸ total CFU/gallon) of the following bacteria:

Bacillus subtilis 5.30x10⁴ CFU/ml
Bacillus methylotrophicus . . 5.30x10⁴ CFU/ml

Microorganisms exempt from CFR requirements – 40 CFR 725.
Store in a cool, dark place to maximize shelf life.
Expires 12 months from delivery date.



Guaranteed by: AgroLiquid
Division of COG Marketers, Ltd.
3055 W M-21
St. Johns, MI 48879
agroliquid.com



Directions For Use General Guideline:

For proper agronomic application rates suitable for your geographical area or the maximum allowable non-nutrient application rate per acre, consult a trained soil specialist at AgroLiquid or call or write to AgroLiquid with the address provided.

PrimAgro K™ is an advanced, high-efficiency potassium solution containing sulfur with a near perfect pH for the effective delivery of organically derived biological components and BioActivites™. The PrimAgro Technology in K deploys targeted microbe populations that decompose organic matter in the root zone releasing potassium while strengthening the potassium absorption mechanisms within the plant. PrimAgro K is also chloride- and hydroxide- free.

Crop	In-Furrow
Corn (Grain)	3-8 gal/A
30" Row Spacing	
Corn (Silage)	3-8 gal/A
30" Row Spacing	
Soybeans	0-3 gal/A
30" Row Spacing	
Soybeans	0-6 gal/A
15" Row Spacing	
Sorghum	0-3 gal/A
Dry Beans	0-1 gal/A
Cotton	0-3 gal/A
Sugarbeet	0-1 gal/A
Canola	0-5 gal/A
Drilled or air-seeded	
Wheat	2-10 gal/A
(Spring or Winter)	
Potato	3-5 gal/A
	Direct contact with the seed piece

Broadcast, or banded not less than 2" from the seed furrow, surface banded, or applied through drip irrigation at the base of the plant

RATE: 1-20 gal/A

Corn	Wheat	Potato
Canola	Alfalfa	

RATE: 1-15 gal/A

Soybean	Sorghum
---------	---------

RATE: 1-10 gal/A

Dry Beans	Sugarbeet
Cotton	

RATE: 3-5 gal/A

Grapes

In-Season Soil Application

Use Rate Per Application

Corn

Sidedress; RATE: 1-20 gal/A

Soybean 30" Rows

Sidedress; RATE: 1-15 gal/A

Sorghum

Sidedress; RATE: 1-15 gal/A

Dry Beans

Sidedress; RATE: 1-10 gal/A

Cotton

Sidedress; RATE: 1-10 gal/A

Sugarbeet

Sidedress; RATE: 1-10 gal/A

Wheat

Topdress up to growth stage Feekes 4; RATE: 3-5 gal/A

Potato

Sidedress or fertigation; RATE: 1-10 gal/A

Alfalfa

Broadcast prior to or within 14 days after spring green-up and/or 0-7 days after cutting; RATE 1-10 gal/A

Grapes

Broadcast, surface banded or through drip irrigation at bud break or during the growing season; RATE: 3-5 gal/A

Tomato

Banded or through drip irrigation during the growing season; RATE: 3-5 gal/A

Tobacco

Banded or through drip irrigation during the growing season; RATE: 3-7 gal/A

Apples

Banded or through drip irrigation during the growing season; RATE: 3-8 gal/A

Tree Nuts

Banded or through drip irrigation during the growing season; RATE: 3-8 gal/A

Other Tree Fruits

Banded or through drip irrigation during the growing season; RATE: 3-8 gal/A

Vegetables

Broadcast, surface banded or through drip irrigation during the growing season; RATE: 3-5 gal/A

Please consult with an AgroLiquid Sales Account Manager or Agronomist for further direction when utilizing rates higher than the lower limit of the given range.

